



Case Report

Rare acute idiopathic subdural hematoma: A case report and literature review

Adilson de Oliveira^{1,2}, Wellingson da Silva Paiva¹, Manoel Jacobsen Teixeira¹

¹Department of Neurosurgery, Hospital das Clinicas da Faculdade de Medicina da USP, Sao Paulo, Brazil, ²Neurosurgery Service, Clinica Girassol, Luanda, Angola.

E-mail: *Adilson de Oliveira - adilsonvalmont@gmail.com; Wellingson da Silva Paiva - wellingsonpaiva@yahoo.com.br; Manoel Jacobsen Teixeira - manoeljacobson@gmail.com



***Corresponding author:**

Adilson de Oliveira,
Department of Neurosurgery,
Faculty of Medicine, São Paulo
University, São Paulo, Brazil;
Serviço de Neurocirurgia Clínica
Girassol, Luanda, Angola.

adilsonvalmont@gmail.com

Received : 21 December 19

Accepted : 30 December 19

Published : 17 January 20

DOI:

10.25259/SNI_499_2019

Quick Response Code:



ABSTRACT

Background: Acute spontaneous subdural hematoma is rare. For patients under 40 years of age, we found only five previous reports. Here, we have presented a sixth case study.

Case Description: A 27-year-old male initially presented with a high-intensity headache without any neurological deficits. The brain computed tomography revealed a left frontoparietal lesion, consistent with an acute epidural hematoma. However, the bone window examination showed no fracture, and at surgery, this lesion proved to be an acute subdural hematoma. Additional studies, including cerebral angiography, brain magnetic resonance imaging, and a complete coagulation work-up, were all negative.

Conclusion: This case report and literature review focused on the rarity of acute idiopathic/spontaneous subdural hematomas.

Keywords: Intracranial hematoma, Neurosurgery, Subdural hematoma, Traumatic brain injury

INTRODUCTION

There are few documented cases of acute spontaneous subdural hematomas (ASSDH) occurring in healthy young men without a history of trauma [Table 1].^[4] Here, we present a 27-year-old male with an ASSDH and reviewed five other cases of idiopathic ASSDH in the patient under 40 years of age.

CASE REPORT

A 27-year-old male presented with a high-intensity headache of 3 h duration. He exhibited no focal neurological deficit or any laboratory/coagulation abnormalities [Table 2]. The brain computed tomography (CT) scan documented a left frontoparietal lesion (e.g., 16 mm side), consistent with an acute epidural hematoma. However, the bone window CT showed no underlying fracture, and at surgery (e.g., a routine craniotomy), the lesion proved to be an acute subdural hematoma [Figure 1]. The postoperative CT confirmed adequate removal of the clot [Figure 2]. Subsequently, the patient's additional studies including cerebral angiography, brain magnetic resonance (MR), and an additional full coagulation work-up all proved negative [Figures 2-5].

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

©2020 Published by Scientific Scholar on behalf of Surgical Neurology International




Figure 1: The preoperative computed tomography (CT) scan showed the left parietal subdural hematoma. Due to its lenticular configuration, this could easily be misinterpreted as an epidural hematoma.




Figure 4: Left cerebral angiography was normal.




Figure 2: Notably, the bone window CT demonstrated no accompanying skull fracture.




Figure 5: Postoperative T1-weighted enhanced brain magnetic resonance (MR) imaging. The postoperative brain MR performed with gadolinium diethylenetriaminepentaacetic acid was negative.



Figure 3: The postoperative computed tomography scan showed complete resection of the hematoma.

DISCUSSION

History of ASSDH

Munro, in 1934, was the first to report an ASSDH; a decade later, Scott reported two more cases.^[9] In 1971, Talalla and McKissock coined the phrase “acute spontaneous SDH (ASSDH).”^[10,11] The previous reports indicated that ASSDH typically occurred in male teenagers and correlated with good outcomes when diagnosed and treated early in the clinical course (e.g., before the onset of a severe neurological deficit).^[2,5,8,13]

Risk factors for ASSDH

Risk factors for ASSDH included hypertension, vascular malformations, neoplasia (e.g., hematological malignancies causing thrombocytopenia), other solid tumors, dural metastases, hypervitaminosis, coagulopathy/alcoholism, and

Table 1: Spontaneous subdural hematoma in patients under 40 years old reported in literature.

Reference author/year	Age M/F	CT/MR	Surgery	Outcome
Kulali, 1992	15 M	Right temporoparietal subdural hematoma with midline shift	No surgical treatment	Normal neurological examination
Man Ho, 1994	39 F	Left frontoparietal subdural hematoma with midline shift	Craniotomy and hematoma evacuation	Hemiparesis
Arnold <i>et al.</i> , 2011 ^[1]	15 M	Left frontoparietal subdural hematoma with midline shift	Craniotomy and hematoma evacuation	Normal neurological examination
Arnold <i>et al.</i> , 2011 ^[1]	16 M	Right frontoparietal subdural hematoma with midline shift	Craniotomy and hematoma evacuation	Normal neurological examination
Brennan and Fuller, 2011 ^[3]	37 M	Right frontoparietal subdural hematoma 28 mm in depth, with 13 mm of midline shift	Craniotomy and hematoma evacuation	Normal neurological examination
de Oliveira, 2019*	27 M	16 mm thick acute left frontoparietal extradural hematoma	Craniotomy, intraoperative find of subdural hematoma and evacuation	Normal neurological examination

*Our case; M: Male, F: Female, CT: Computed tomography, MR: Magnetic resonance

Table 2: Laboratorial investigation.

Exam	Value	Normal range
Hemoglobin	17.9 g/dl	13–18 g/dl
Partial thromboplastin time	25.4 s	20.8–31.2 s
Prothrombin time	11.6 s	Control 11.6 s
INR	1.0	0.95–1.2
R	0.98	0.80–1.2
Platelet counts	201.000/mm ³	140.000–450.000/mm ³
Antithrombin	109%	79–131%
Factor VIII	177%	50–150%
Factor IX	124%	60–150%

bleeding from cerebral artery aneurysms/cortical arteries. In the case presented, the patient had none of these risk factors. Notable, however, was the CT finding of a lenticular clot pathognomonic for an epidural hematoma (e.g., only 8% of subdural hematomas demonstrate this radiological shape), but without a fracture on the bone window CT.^[12] At surgery, this proved to be an ASSDH.

CONCLUSION

Spontaneous intracranial hematomas are rare life-threatening lesions that typically present with mild symptoms and less severe neurological findings versus traumatic acute subdural hematomas. In addition to obtaining preoperative noncontrast CT bone and soft-tissue studies, patients postoperatively should undergo brain MR scans, cerebral angiography, and a full coagulation work-up to rule out other etiologies of these rare lesions.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Arnold PM, Christiano LD, Klemp JA, Anderson KK. Nontraumatic spontaneous acute subdural hematoma in identical teenage twins 1 year apart. *Pediatr Emerg Care* 2011;27:649-51.
2. Avis SP. Nontraumatic acute subdural hematoma. A case report and review of the literature. *Am J Forensic Med Pathol* 1993;14:130-4.
3. Brennan PM, Fuller E, Shanmuganathan M, Keston P, Fouyas I. Spontaneous subdural haematoma in a healthy young male. *BMJ Case Rep* 2011;2011:bcr0120113694.
4. Coombs JB, Coombs BL, Chin EJ. Acute spontaneous subdural hematoma in middle aged adult: Case report and literature review. *J Emerg Med* 2014;212:1-6.
5. Depreitere B, Van Calenbergh F, van Loon J. A clinical comparison of non-traumatic acute subdural haematomas either related to coagulopathy or of arterial origin without coagulopathy. *Acta Neurochir (Wien)* 2003;145:541-6.
6. Kulah A, Taşdemir N, Fiskeci C. Acute spontaneous subdural hematoma in a teenager. *Childs Nerv Syst* 1992;8:343-6.
7. Marconi F, Fiori L, Parenti G, Ravelli V. Acute spontaneous subdural haematoma. Description of four clinical cases. *J Neurosurg Sci* 1991;35:97-102.
8. Missori P, Fenga L, Maraglino C, Rocchi G, Nardacci B, Calderaro G, *et al.* Spontaneous acute subdural hematomas. A clinical comparison with traumatic acute subdural hematomas. *Acta Neurochir (Wien)* 2000;142:697-701.
9. Monsivais D, Choi HA, Kitagawa R, Franch M,

- Cai C. A retrospective analysis of surgical outcomes for acute subdural hematoma in an elderly cohort. *Interdiscip Neurosurg* 2018;14:130-4.
10. Munro D. The diagnosis and treatment of subdural hematoma. A report of sixty two cases. *N Engl J Med* 1934;210:1145-60.
11. Scott M. Spontaneous nontraumatic subdural hematomas. *J Am Med Assoc* 1949;141:596-602.
12. Su IC, Wang KC, Huang SH, Li CH, Kuo LT, Lee JE, *et al.* Differential CT features of acute lentiform subdural hematoma and epidural hematoma. *Clin Neurol Neurosurg* 2010;112:552-6.
13. Talalla A, McKissock W. Acute “spontaneous” subdural hemorrhage. An unusual form of cerebrovascular accident. *Neurology* 1971;21:19-25.

How to cite this article: de Oliveira A, da Silva Paiva W, Teixeira MJ. Rare acute idiopathic subdural hematoma: A case report and literature review. *Surg Neurol Int* 2020;11:9.